

**PDC2019
Washington, DC, USA**

- Key International and Political Developments**
- Advancements and Progress in NEO Discovery**
- NEO Characterization Results**
- Deflection and Disruption Models & Testing**
- Mission & Campaign Designs**
- Impact Consequences**
- Disaster Response**
- Decision to Act**
- Public Education & Communication**

“EFFECTIVE HEIGHT OF BURST” REVISTED

Mark Boslough⁽¹⁾ and Amanda R. Truitt⁽²⁾

Los Alamos National Laboratory, P.O. Box 1663, Los Alamos, NM 87545, USA

⁽¹⁾ *Mail Stop T087, 505-999-7756, mbeb@unm.edu*

⁽²⁾ *Mail Stop T087, 505-664-0035, atruitt@lanl.gov*

Keywords: *Tunguska, airburst, hydrocode, modelling, consequences*

ABSTRACT

In 1993, Chyba et al. used a pancake model to argue that the 1908 Tunguska airburst was due to a stony asteroid. To estimate the effects at the surface, they used the altitude of maximum energy deposition as a proxy for the equivalent static height of burst and used the scaling laws from nuclear weapons effects literature. After the 1994 impact of Comet Shoemaker-Levy 9, different modeling groups came to different conclusions about the maximum depth achieved by fragments, with the Sandia group arguing for deeper penetration due in part to continued downward flow of vaporized comet after energy deposition was complete. At the 2007 Planetary Defense Conference, Boslough made a similar case for Tunguska, arguing that the “effective height of burst” can be significantly lower than the altitude of maximum energy deposition and concluding that the observed damage can be attributed to downward-directed airburst with a lower total kinetic yield (3-5 Mt as opposed to 10-20 Mt). Recent work by Wheeler & Mathias (2019, in press) rejects the notion that the effective height of burst is lower than the peak energy deposition point, and concludes (in part on that basis) that the most probable Tunguska-scale yield is 20-30 Mt. Because a Tunguska-scale airburst is by far the most likely NEO disaster to take place in our lifetimes, risk assessments rely heavily on our estimates of damage as a function of size and frequency. More certainty in our quantitative risk assessments will require that we converge on a consensus about the size of the Tunguska event, which in turn requires that we come to an agreement on the difference (if any) between “effective height of burst” and peak energy deposition altitude. To this end, we will present new CTH and RAGE simulations in which we isolate that effect as a function of a wide range of impactor mass, entry angle, and velocity.

Comments:

This work was supported by the US Department of Energy through the Los Alamos National Laboratory. Los Alamos National Laboratory is operated by Triad National Security, LLC, for the National Nuclear Security Administration of U.S. Department of Energy (Contract No. 89233218CNA000001).